arXiv:0705.0627v1 [math-ph] 4 May 2007

# The Intergals of Motion for the Deformed $W$-Algebra $W_{q,t}(\widehat{sl_N})$

Boris FEIGIN [α], Takeo KOJIMA [β], Jun'ichi SHIRAISHI [γ], Hidekazu WATANABE [γ]

[α] *L.D.Landau Institute for Theoretical Sciences,*
*Chernogolovka, Moscow 142432, RUSSIA*
[β] *Department of Mathematics, College of Science and Technology, Nihon University,*
*Surugadai, Chiyoda-ku, Tokyo 101-0062, JAPAN*
[γ] *Graduate School of Mathematical Science, University of Tokyo,*
*Komaba, Meguro-ku, Tokyo, 153-8914, JAPAN*

## Abstract

We review the deformed $W$-algebra $W_{q,t}(\widehat{sl_N})$ and its screening currents. We explicitly construct the Local Integrals of Motion $\mathcal{I}_n$ $(n = 1, 2, \cdots)$ for this deformed $W$-algebra. We explicitly construct the Nonlocal Integrals of Motion $\mathcal{G}_n$ $(n = 1, 2, \cdots)$ by means of the screening currents. Our Integrals of Motion commute with each other, and give the elliptic version of those for the Virasoro algebra and the $W$-algebra $W(\widehat{sl_3})$ [1, 2].

# 1 Introduction

V.Bazhanov, S.Lukyanov and Al.Zamolodchikov [1] constructed field theoretical analogue of the commuting transfer matrix $\mathbf{T}(z)$ as the trace of the free field realization of the monodromy matrix associated with the quantum affine symmetry $U_q(\widehat{sl_2})$. They call the coefficients of the asymptotic expansion of the operator $\log\mathbf{T}(z)$ at $z \to \infty$, the Local Integrals of Motioin for the Virasoro algebra. The Local Integrals of Motion $I_n$ have the form

$$I_{2k-1} = \int_0^{2\pi} \frac{du}{2\pi} T_{2k}(u), \quad (k = 1, 2, \cdots). \tag{1.1}$$

Here the densities $T_{2k}(u)$ are differential polynomials of the energy momentum tensor of the Virasoro algebra $T(u) = -\frac{C_{CFT}}{24} + \sum_{n=-\infty}^{\infty} L_{-n} e^{\sqrt{-1}nu}$, where the operators $L_n$ satisfies the commutation relation, $[L_n, L_m] = (n-m)L_{n+m} + \frac{C_{CFT}}{12}(n^3 - n)\delta_{n+m,0}$. The first few densities $T_{2k}(u)$ are written by

$$T_2(u) = T(u),\ T_4(u) =: T^2(u) :,\ T_6(u) =: T^3(u) : + \frac{C_{CFT}+2}{12} : (T'(u))^2 : . \tag{1.2}$$

However they did not give explicit formulae of general densities $T_{2k}$, all densities $T_{2k}(u)$ are uniequely determined by requirement of the commutativity, $[I_{2k-1}, I_{2l-1}] = 0,\ (k, l = 1, 2, \cdots)$. They call the coefficients of the Taylor expnsion of the operator $\mathbf{T}(z)$ at $z = 0$, the Nonlocal Integrals of Motion. They gave explicit formulae for the Nonlocal Integrals of Motion $G_k,\ (k = 1, 2, \cdots)$, by means of the screening currents $F_1(z), F_2(z)$.

$$\begin{aligned} G_k &= \int \cdots \int_{2\pi \geqq u_1 \geqq u_2 \geqq \cdots \geqq u_{2k} \geqq 0} (e^{2\pi\sqrt{-1}P} F_1(u_1)F_2(u_2)F_1(u_3)F_2(u_4)\cdots F_2(u_{2k}) \\ &+ e^{-2\pi\sqrt{-1}P} F_2(u_1)F_1(u_2)F_2(u_3)F_1(u_4)\cdots F_1(u_{2k}))du_1du_2\cdots du_{2k}, \end{aligned} \tag{1.3}$$

where $P$ is zero-mode operator. Because of the commutativity of the transfer matrix $\mathbf{T}(z)$, the integtrals of motion commute with each other : $[I_{2k-1}, I_{2l-1}] = [I_{2k-1}, G_l] = [G_k, G_l] = 0$. V.Bazhanov, A.Hibberd and S.Khoroshkin [2] considered the $W$-algebra $W(\widehat{sl_3})$ version of [1].

V.Bazhanov, S.Lukyanov and Al. Zamolodchikov's theory [1] can be regarded as the quantum version of the KdV problem, as it reduces to the classical KdV problem in classical limit $C_{CFT} \to -\infty$, under the substitution

$$T(u) \to -\frac{C_{CFT}}{6} U(u), \quad [\ ,\ ] \to \frac{6\pi}{\sqrt{-1}C_{CFT}} \{\ ,\ \}. \tag{1.4}$$

The Poisson bracket structure $\{,\}$ gives the second Hamiltonian structure of the KdV equation. The Local Integrals of Motion $I_{2k-1}$ tend to $I_{2k-1}^{(cl)}$.

$$I_{2k-1}^{(cl)}=\int_0^{2\pi}\frac{du}{2\pi}T_{2k}^{(cl)}(u),\quad (k=1,2,\cdots),\tag{1.5}$$

where the first few densities are written by

$$T_2^{(cl)}=U(u),\quad T_4^{(cl)}=U^2(u),\quad T_6^{(cl)}=U^3(u)-\frac{1}{2}(U'(u))^2.\tag{1.6}$$

The KdV hierarchy are given by

$$\partial_{t_{2k-1}}U=\{I_{2k-1}^{(cl)},U\},\ (k=1,2,\cdots).\tag{1.7}$$

The purpose of this paper is to construct the elliptic version of the Integrals of Motion $I_k$ and $G_k$ given by V.Bazhanov, S.Lukyanov and Al.Zamolodchikov [1] and V.Bazhanov, A.Hibberd and S.Khoroshkin [2]. Our method of construction is completely different from those of [1, 2]. Instead of considering the transfer matrix $\mathbf{T}(z)$, we directly give explicit formulae of both Local Integrals of Motion $\mathcal{I}_n$ and the Nonlocal Integrals of Motion $\mathcal{G}_m$, for the deformed $W$-algebra $W_{q,t}(\widehat{sl_N})$.

$$[\mathcal{I}_m,\mathcal{I}_n]=0,\quad [\mathcal{G}_m,\mathcal{G}_n]=0,\quad [\mathcal{I}_m,\mathcal{G}_n]=0,\quad (m,n=1,2,\cdots).\tag{1.8}$$

The organization of this paper is as follows. In Section 2, we give basic definition, including bosons, screening currents. In Section 3, we review the deformed $W$-algebra $W_{q,t}(\widehat{sl_N})$. In Section 4, we construct explicit formulae for the Local Integrals of Motion $\mathcal{I}_n$, $(n=1,2,\cdots)$ for the deformed $W$-algebra $W_{q,t}(\widehat{sl_N})$. In Section 5, we construct explicit formulae for the Nonlocal Integrals of Motion $\mathcal{G}_m$, $(m=1,2,\cdots)$.

# 2 Basic Definition

In this section we give the basic definition. Let us fix three parameters $0<x<1, r\in\mathbb{C}$ and $s\in\mathbb{C}$.

## 2.1 Bosons

Let $\epsilon_i (1\leqq i\leqq N)$ be an orthonormal basis in $\mathbb{R}^N$ relative to the standard basis in $\mathbb{R}^N$ relative to the standard inner product $(,)$. Let us set $\bar{\epsilon}_i=\epsilon_i-\epsilon, \epsilon=\frac{1}{N}\sum_{j=1}^N\epsilon_j$. We idetify $\epsilon_{N+1}=\epsilon_1$. Let $P=\sum_{i=1}^N\mathbb{Z}\bar{\epsilon}_i$ the weight lattice. Let us set $\alpha_i=\bar{\epsilon}_i-\bar{\epsilon}_{i+1}\in P$.

Let $\beta_m^j$ be the oscillators ($1 \leqq j \leqq N, m \in \mathbb{Z} - \{0\}$) with the commutation relations

$$[\beta_m^i, \beta_n^j] = \begin{cases} m\frac{[(r-1)m]}{[rm]}\frac{[(s-1)m]}{[sm]}\delta_{n+m,0} & (1 \leqq i = j \leqq N) \\ -m\frac{[(r-1)m]}{[rm]}\frac{[m]}{[sm]}x^{sm\ \mathrm{sgn}(i-j)}\delta_{n+m,0} & (1 \leqq i \neq j \leqq N) \end{cases} \tag{2.1}$$

Here the symbol $[a]$ stands for $\frac{x^a - x^{-a}}{x - x^{-1}}$.

We also introduce the zero mode operator $P_\lambda$, $(\lambda \in P)$. They are $\mathbb{Z}$-linear in $\lambda$ and satisfy

$$[iP_\lambda, Q_\mu] = (\lambda, \mu), \quad (\lambda, \mu \in P). \tag{2.2}$$

Let us intrduce the bosonic Fock space $\mathcal{F}_{l,k}(l, k \in P)$ generated by $\beta_{-m}^j(m > 0)$ over the vacuum vector $|l, k\rangle$ :

$$\mathcal{F}_{l,k} = \mathbb{C}[\{\beta_{-1}^j, \beta_{-2}^j, \cdots\}_{1 \leqq j \leqq N}]|l, k\rangle, \tag{2.3}$$

where

$$\beta_m^j|l, k\rangle = 0, (m > 0), \tag{2.4}$$

$$P_\alpha|l, k\rangle = \left(\alpha, \sqrt{\frac{r}{r-1}}l - \sqrt{\frac{r-1}{r}}k\right)|l, k\rangle, \tag{2.5}$$

$$|l, k\rangle = e^{i\sqrt{\frac{r}{r-1}}Q_l - i\sqrt{\frac{r-1}{r}}Q_k}|0, 0\rangle. \tag{2.6}$$

Let us set the Dynkin-diagram automorphism $\eta$ by

$$\eta(\beta_m^1) = x^{-\frac{2s}{N}m}\beta_m^2, \cdots, \eta(\beta_m^{N-1}) = x^{-\frac{2s}{N}m}\beta_m^N,\ \eta(\beta_m^N) = x^{\frac{2s}{N}(N-1)m}\beta_m^1, \tag{2.7}$$

and $\eta(\epsilon_i) = \epsilon_{i+1}$, $(1 \leqq i \leqq N)$.

## 2.2 Basic Operators

In this section we introduce the basic operators. Let us set $z = x^{2u}$.

**Definition 2.1** *We set the screening currents $F_j(z)(1 \leqq j \leqq N)$ by*

$$F_j(z) = e^{i\sqrt{\frac{r-1}{r}}Q_{\alpha_j}}(x^{(\frac{2s}{N}-1)j}z)^{\sqrt{\frac{r-1}{r}}P_{\alpha_j} + \frac{r-1}{r}} \times :\exp\left(\sum_{m \neq 0}\frac{1}{m}B_m^j z^{-m}\right):,\ (1 \leqq j \leqq N-1) \tag{2.8}$$

$$F_N(z) = e^{i\sqrt{\frac{r-1}{r}}Q_{\alpha_N}}(x^{2s-N}z)^{\sqrt{\frac{r-1}{r}}P_{\bar{\epsilon}_N} + \frac{r-1}{2r}}(z)^{-\sqrt{\frac{r-1}{r}}P_{\bar{\epsilon}_1} + \frac{r-1}{2r}} \times :\exp\left(\sum_{m \neq 0}\frac{1}{m}B_m^N z^{-m}\right): . \tag{2.9}$$

*Here we set*

$$B_m^j = (\beta_m^j - \beta_m^{j+1}) x^{-\frac{2s}{N}jm}, \ (1 \leqq j \leqq N-1), \tag{2.10}$$

$$B_m^N = (x^{-2sm}\beta_m^N - \beta_m^1). \tag{2.11}$$

The screening currents $F_j(z)$, $(1 \leqq j \leqq N-1)$ are studied well in [7]. We introduce new current $F_N(z)$, which can be regarded as "affinization" of screenings $F_j(z)$, $(1 \leqq j \leqq N-1)$.

In what follows, the symbol $[u]_r$ stands for the theta function satisfying

$$[u+r]_r = -[u]_r = [-u]_r, \tag{2.12}$$

$$[u+\tau]_r = -e^{\frac{2\pi i}{r}(u+\frac{\tau}{2})}[u]_r, \quad \text{where } \tau = \frac{\pi i}{\log x}. \tag{2.13}$$

Explicitly it is given by

$$[u]_r = x^{u^2/r-u}\Theta_{x^{2r}}(x^{2u}), \tag{2.14}$$

$$\Theta_q(z) = (z;q)_\infty (q/z;q)_\infty (q;q)_\infty, \tag{2.15}$$

$$(z;q)_\infty = \prod_{j=0}^{\infty}(1-zq^j). \tag{2.16}$$

**Proposition 2.1** *The screening currents $F_j(z)$, $(1 \leqq j \leqq N)$ satisfy the following commutation relations,*

$$\frac{1}{[u_1-u_2-\frac{s}{N}+1]_r}F_j(z_1)F_{j+1}(z_2) = \frac{1}{[u_2-u_1+\frac{s}{N}]_r}F_{j+1}(z_2)F_j(z_1), \quad (1 \leqq j \leqq N), \tag{2.17}$$

$$\frac{[u_1-u_2]_r}{[u_1-u_2-1]_r}F_j(z_1)F_j(z_2) = \frac{[u_2-u_1]_r}{[u_2-u_1-1]_r}F_j(z_2)F_j(z_1), \quad (1 \leqq j \leqq N), \tag{2.18}$$

*and*

$$F_i(z_1)F_j(z_2) = F_j(z_2)F_i(z_1), \quad (|i-j| \geqq 2). \tag{2.19}$$

*We read $F_{N+1}(z) = F_1(z)$.*

**Proposition 2.2** *The action of $\eta$ on the screenings $F_j(z)$ is given by*

$$\eta(F_j(z)) = F_{j+1}(x^{1-\frac{2s}{N}}z), \quad (1 \leqq j \leqq N-2), \tag{2.20}$$

$$\eta(F_{N-1}(z)) = F_N(x^{1-\frac{2s}{N}}z)x^{(2s-N)(-\sqrt{\frac{r-1}{r}}P_{\bar{\epsilon}_1}+\frac{r-1}{r})}, \tag{2.21}$$

$$\eta(F_N(z)) = F_1(x^{1-\frac{2s}{N}}z)x^{(2s-N)(\sqrt{\frac{r-1}{r}}P_{\bar{\epsilon}_1}+\frac{r-1}{r})}. \tag{2.22}$$

*We have $\eta(F_{N-1}(z_1)F_N(z_2)) = F_N(z_1)F_1(z_2)$.*

**Definition 2.2** *We set the fundamental operator* $\Lambda_j(z), (1 \leqq j \leqq N)$ *by*

$$\Lambda_j(z) \quad = \quad x^{-2\sqrt{r(r-1)}P_{\bar{\epsilon}_j}} : \exp\left(\sum_{m\neq 0} \frac{x^{rm}-x^{-rm}}{m}\beta_m^j z^{-m}\right) : \quad (1 \leqq j \leqq N). \qquad (2.23)$$

**Proposition 2.3** *The action of* $\eta$ *on the screenings* $\Lambda_j(z)$ *is given by*

$$\eta(\Lambda_j(z)) = \Lambda_{j+1}(xz), \quad (1 \leqq j \leqq N-1), \quad \eta(\Lambda_N(z)) = \Lambda_1(x^{1-2s}z). \qquad (2.24)$$

**Proposition 2.4** *The screening currents* $F_j(z)$, $(1 \leqq j \leqq N)$ *and the fundamental operators* $\Lambda_j(z)$, $(1 \leqq j \leqq N)$ *commute up to delta-function* $\delta(z) = \sum_{n\in\mathbb{Z}} z^m$.

$$[\Lambda_j(z_1), F_j(z_2)] \quad = \quad (-1 + x^{-2r+2})\delta\left(x^{\frac{2s}{N}j-r}\frac{z_2}{z_1}\right) : \Lambda_j(z_1)F_j(z_2) :, \quad (1 \leqq j \leqq N), \qquad (2.25)$$

$$[\Lambda_{j+1}(z_1), F_j(z_2)] \quad = \quad (1 - x^{-2r+2})\delta\left(x^{\frac{2s}{N}j+r}\frac{z_2}{z_1}\right) : \Lambda_{j+1}(z_1)F_j(z_2) :, \quad (1 \leqq j \leqq N-1), \qquad (2.26)$$

$$[\Lambda_1(z_1), F_N(z_2)] \quad = \quad (1 - x^{-2r+2})\delta\left(x^{r}\frac{z_2}{z_1}\right) : \Lambda_1(z_1)F_N(z_2) : . \qquad (2.27)$$

# 3 Deformed $W$-Algebra $W_{q,t}(\widehat{sl_N})$

In this section we review the deformed $W$-algebra $W_{q,t}(\widehat{sl_N})$ [3, 4, 5].

## 3.1 Deformation of $W$-Algebra

In this section we give the deformation of the $W$-algebra.

**Definition 3.1** *Let us set the operator* $T_j(z)$, $(1 \leqq j \leqq N)$ *by*

$$T_j(z) = \sum_{1\leqq s_1 < s_2 < \cdots < s_j \leqq N} : \Lambda_{s_1}(x^{-j+1}z)\Lambda_{s_2}(x^{-j+3}z)\cdots\Lambda_{s_j}(x^{j-1}z) : . \qquad (3.1)$$

**Proposition 3.1** *The bosonic operators* $T_j(z)$, $(1 \leqq j \leqq N)$ *satisfy the following relations.*

$$\begin{aligned} & f_{i,j}(z_2/z_1)T_i(z_1)T_j(z_2) - f_{j,i}(z_1/z_2)T_j(z_2)T_i(z_1) \\ = \quad & c\sum_{k=1}^{i}\prod_{l=1}^{k-1}\Delta(x^{2l+1}) \times \left(\delta\left(\frac{x^{j-i+2k}z_2}{z_1}\right) f_{i-k,j+k}(x^{-j+i})T_{i-k}(x^{-k}z_1)T_{j+k}(x^k z_2)\right. \\ - \quad & \left.\delta\left(\frac{x^{-j+i-2k}z_2}{z_1}\right) f_{i-k,j+k}(x^{j-i})T_{i-k}(x^{k}z_1)T_{j+k}(x^{-k}z_2)\right), \quad (1 \leqq i \leqq j \leqq N), (3.2) \end{aligned}$$

*where* $\delta(z)=\sum_{n\in\mathbb{Z}}z^n$.

*Here we set the constnt* $c$ *and the auxiliary function* $\Delta(z)$ *by*

$$c=-\frac{(1-x^{2r})(1-x^{-2r+2})}{(1-x^2)},\quad \Delta(z)=\frac{(1-x^{2r-1}z)(1-x^{1-2r}z)}{(1-xz)(1-x^{-1}z)}. \tag{3.3}$$

*Here we set the structure functions,*

$$f_{i,j}(z)=\exp\left(\sum_{m=1}^{\infty}\frac{1}{m}(1-x^{2rm})(1-x^{-2(r-1)m})\frac{(1-x^{2mMin(i,j)})(1-x^{2m(s-Max(i,j))})}{(1-x^{2m})(1-x^{2sm})}x^{|i-j|m}z^m\right). \tag{3.4}$$

**Example** For $N=2$ the operators $T_1(z),T_2(z)$ satisfy

$$\begin{aligned}&f_{1,1}(z_2/z_1)T_1(z_1)T_1(z_2)-f_{1,1}(z_1/z_2)T_1(z_2)T_1(z_1)\\ =\ &c(\delta(x^2z_2/z_1)T_2(xz_2)-\delta(x^2z_1/z_2)T_2(x^{-1}z_2)), &(3.5)\\ &f_{1,2}(z_2/z_1)T_1(z_1)T_2(z_2)=f_{2,1}(z_1/z_2)T_2(z_2)T_1(z_1), &(3.6)\\ &f_{2,2}(z_2/z_1)T_2(z_1)T_2(z_2)=f_{2,2}(z_1/z_2)T_2(z_2)T_2(z_1). &(3.7)\end{aligned}$$

**Example** For $N=3$ the operators $T_1(z),T_2(z),T_3(z)$ satisfy

$$\begin{aligned}&f_{1,1}(z_2/z_1)T_1(z_1)T_1(z_2)-f_{1,1}(z_1/z_2)T_1(z_2)T_1(z_1)\\ =\ &c(\delta(x^2z_2/z_1)T_2(xz_2)-\delta(x^2z_1/z_2)T_2(x^{-1}z_2)), &(3.8)\\ &f_{1,2}(z_2/z_1)T_1(z_1)T_2(z_2)-f_{2,1}(z_1/z_2)T_2(z_2)T_1(z_1)\\ =\ &c(\delta(x^3z_2/z_1)T_3(xz_2)-\delta(x^3z_1/z_2)T_3(x^{-1}z_2)), &(3.9)\\ &f_{2,2}(z_2/z_1)T_2(z_1)T_2(z_2)-f_{2,2}(z_1/z_2)T_2(z_2)T_2(z_1)\\ =\ &cf_{1,3}(1)(\delta(x^2z_2/z_1)T_1(xz_2)T_3(xz_2)-\delta(x^2z_1/z_2)T_1(x^{-1}z_2)T_3(x^{-1}z_2)), &(3.10)\\ &f_{1,3}(z_2/z_1)T_1(z_1)T_3(z_2)=f_{3,1}(z_1/z_2)T_3(z_2)T_1(z_1), &(3.11)\\ &f_{2,3}(z_2/z_1)T_2(z_1)T_3(z_2)=f_{3,2}(z_1/z_2)T_3(z_2)T_2(z_1), &(3.12)\\ &f_{3,3}(z_2/z_1)T_3(z_1)T_3(z_2)=f_{3,3}(z_1/z_2)T_3(z_2)T_3(z_1). &(3.13)\end{aligned}$$

**Definition 3.2** *The three parameter deformed* $W$*-algebra is an associative algebra generated by* $T_n^{(j)}$, $(n\in\mathbb{Z},1\leqq j\leqq N)$. *Defining relations are given by (3.2). The elements* $T_n^{(j)}$ *are Fourier coefficients of* $T_j(z)=\sum_{n\in\mathbb{Z}}T_n^{(j)}z^{-n}$.

For general $N$, upon specialization $s=N$, the operator $T_N(z)$ degenerate to scalar 1. Let us set parameters $q=x^{2r}$ and $t=x^{2r-2}$.

**Theorem 3.2** *Upon specialization* $s = N$*, the operator* $T_j(z)$ $(1 \leqq j \leqq N)$ *give a free field realization of the deformed* $W$*-algebra* $W_{q,t}(\widehat{sl_N})$*.*

**Example** For $N = 2$ and $s = 2$, the operator $T_2(z)$ degenerates to scalar 1, and Fourier coefficients of $T_1(z) = \sum_{n\in\mathbb{Z}} T_n z^{-n}$ satisfy the defining relation of the deformed Virasoro algebra $Vir_{q,t} = W_{q,t}(\widehat{sl_2})$ [3].

$$[T_n, T_m] = -\sum_{l=0}^{\infty} f_l (T_{n-l}T_{m+l} - T_{m-l}T_{n+l}) + c((q/t)^n - (t/q)^n)\delta_{n+m,0}, \tag{3.14}$$

where we set the structure constant $f_l$ by $f_{11}(z) = 1 + \sum_{l=1}^{\infty} f_l z^l$. Let us set $q = e^h$ and $t = q^\beta$, $(\beta = \frac{r-1}{r})$, and take the limit $h \to 0$ under the following $h$-expansion,

$$T_n = 2\delta_{n,0} + \beta\left(L_n + \frac{(1-\beta)^2}{4\beta}\delta_{n,0}\right)h^2 + O(h^4), \tag{3.15}$$

we have the defining relation of the Virasoro algebra,

$$[L_m, L_n] = (n-m)L_{m+n} + \frac{C_{CFT}}{12}n(n^2-1)\delta_{n+m,0}, \quad C_{CFT} = 1 - \frac{6(1-\beta)^2}{\beta}. \tag{3.16}$$

## 3.2 Comparsion with another definition

At first glance, our definition of the deformed $W$-algebra is different from those in [4, 5]. In this section we show they are essentially the same thing. Let us set the element $\mathcal{C}_m$ by

$$\mathcal{C}_m = \sum_{j=1}^{N} x^{(N-2j+1)m}\beta_m^j. \tag{3.17}$$

This element $\mathcal{C}_m$ is $\eta$-invariant, $\eta(\mathcal{C}_m) = \mathcal{C}_m$. Let us divide $\Lambda_j(z)$ into $\Lambda_j^{DWA}(z)$ and $\mathcal{Z}(z)$.

$$\Lambda_j(z) \quad = \quad \Lambda_j^{DWA}(z)\mathcal{Z}(z), \quad (1 \leqq j \leqq N), \tag{3.18}$$

where we set

$$\Lambda_j^{DWA}(z) \quad = \quad x^{-2\sqrt{r(r-1)}P_{\bar{\epsilon}_j}} : \exp\left(\sum_{m\neq 0}\frac{x^{rm} - x^{-rm}}{m}\left(\beta_m^j - \frac{[m]_x}{[Nm]_x}\mathcal{C}_m\right)z^{-m}\right) : \tag{3.19}$$

$$\mathcal{Z}(z) \quad = \quad : \exp\left(\sum_{m\neq 0}\frac{x^{rm} - x^{-rm}}{m}\frac{[m]_x}{[Nm]_x}\mathcal{C}_m z^{-m}\right) : . \tag{3.20}$$

Let us set

$$T_j^{DWA}(z) = \sum_{1\leqq s_1 < s_2 < \cdots < s_j \leqq N} : \Lambda_{s_1}^{DWA}(x^{-j+1}z)\Lambda_{s_2}^{DWA}(x^{-j+3}z)\cdots\Lambda_{s_j}^{DWA}(x^{j-1}z) : . \tag{3.21}$$

**Theorem 3.3** *The operators* $T_j^{DWA}(z)$, $(1 \leqq j \leqq N-1)$ *give a free field realization of the deformed* $W$*-algebra* $W_{q,t}(\widehat{sl_N})$ *[4, 5].*

**Proposition 3.4** *The operators* $T_j^{DWA}(z)$ *and* $\mathcal{Z}(z)$ *commutes with each other.*

$$T_j^{DWA}(z_1)\mathcal{Z}(z_2) = \mathcal{Z}(z_2)T_j^{DWA}(z_1), \quad (1 \leqq j \leqq N-1). \tag{3.22}$$

Three paramedter deformed $W$-algebra defined in the previous subsection (3.2), can be regarded as an extension of $W_{q,t}(\widehat{sl_N})$, therefore, we sometime call three parameter deformed $W$-algebra, "the deformed $W$-algebra $W_{q,t}(\widehat{sl_N})$". In what follows, mainly, we consider three parameter deformed $W$-algebra. It is simpler to show the commutation relations of the Integrals of Motion $[\mathcal{I}_m, \mathcal{I}_n] = [\mathcal{G}_m, \mathcal{G}_n] = [\mathcal{I}_m, \mathcal{G}_n] = 0$ for three parameter $(x, r, s)$ deformed case than those for two parameter $(x, r, s = N)$ deformed case. One additionnal parameter $s$ resolves singularity in the Integrals of Motion, and make problem simpler.

# 4 Local Integrals of Motion

In this section we give explicit formulae of the Local Integrals of Motion $\mathcal{I}_n$.

In what follows we use the notation of the ordered product.

$$\prod_{\overrightarrow{l \in L}} T_j(z_l) = T_j(z_{l_1})T_j(z_{l_2}) \cdots T_j(z_{l_m}), \quad (L = \{l_1, l_2, \cdots, l_m | l_1 < l_2 < \cdots < l_m\}). \tag{4.1}$$

For formal power series $\mathcal{A}(z_1, z_2, \cdots, z_n) = \sum_{k_1, k_2, \cdots, k_n \in \mathbb{Z}} a_{k_1, k_2, \cdots, k_n} z_1^{k_1} z_2^{k_2} \cdots z_n^{k_n}$, we set the symbol $[\cdots]_{1, z_1, z_2, \cdots, z_n}$.

$$[\mathcal{A}(z_1, z_2, \cdots, z_n)]_{1, z_1, z_2, \cdots, z_n} = a_{0,0,\cdots,0}. \tag{4.2}$$

Let us set the auxiliary function $g_{i,j}(z)$ by fusion of $g_{1,1}(z) = f_{1,1}(z)$.

$$\begin{aligned} g_{i,1}(z) &= g_{1,1}(x^{-i+1}z)g_{1,1}(x^{-i+3}z) \cdots g_{1,1}(x^{i-1}z), \\ g_{i,j}(z) &= g_{i,1}(x^{-j+1}z)g_{i,1}(x^{-j+3}z) \cdots g_{i,1}(x^{j-1}z). \end{aligned} \tag{4.3}$$

**Definition 4.1** *We set the operator* $\mathcal{O}_n(z_1, z_2, \cdots, z_n)$ *by*

$$\mathcal{O}_n(z_1, z_2, \cdots, z_n) = \sum_{\substack{\alpha_1, \alpha_2, \alpha_3, \cdots, \alpha_N \geqq 0 \\ \alpha_1 + 2\alpha_2 + 3\alpha_3 + \cdots + N\alpha_N = n}} \sum_{\substack{A_1^{(1)}, \cdots, A_{\alpha_1}^{(1)}, \cdots, A_1^{(N)}, \cdots, A_{\alpha_N}^{(N)} \subset \{1, 2, \cdots, n\} \\ |A_j^{(t)}| = t, \quad \oplus_{j,t} A_j^{(t)} = \{1, 2, \cdots, n\}}}$$

$$\times \prod_{\overrightarrow{j\in A^{(1)}_{Min}}} T_1(z_j) \prod_{\overrightarrow{j\in A^{(2)}_{Min}}} T_2(x^{-1}z_j) \cdots \prod_{\overrightarrow{j\in A^{(t)}_{Min}}} T_t(x^{-1+t-2[\frac{t}{2}]}z_j) \cdots \prod_{\overrightarrow{j\in A^{(N)}_{Min}}} T_N(x^{-1+N-2[\frac{N}{2}]}z_j)$$

$$\times \prod_{t=1}^{N} \left( (-c)^{t-1} \prod_{u=1}^{t-1} \Delta(x^{2u+1})^{t-u-1} \right)^{\alpha_t} \prod_{t=1}^{N} \prod_{\substack{j=1 \\ j_1=A^{(t)}_{j,1} \\ \cdots \\ j_t=A^{(t)}_{j,t}}}^{\alpha_t} \sum_{\substack{\sigma\in S_t \\ \sigma(1)=1}} \prod_{\substack{u=1 \\ u\neq[\frac{t}{2}]+1}}^{t} \delta\left(\frac{x^2 z_{j_{\sigma(u+1)}}}{z_{j_{\sigma(u)}}}\right)$$

$$\times \prod_{t=1}^{N} \prod_{\substack{j<k \\ j,k\in A^{(t)}_{Min}}} g_{t,t}\left(\frac{z_k}{z_j}\right) \prod_{1\leqq t<u\leqq N} \prod_{\substack{j\in A^{(t)}_{Min} \\ k\in A^{(u)}_{Min}}} g_{t,u}\left(x^{u-t-2[\frac{u}{2}]+2[\frac{t}{2}]}\frac{z_k}{z_j}\right). \tag{4.4}$$

*Here we have set the constant $c$ and the function $\Delta(z)$ in (3.3). When the index set $A^{(t)}_j = \{j_1, j_2, \cdots, j_t | j_1 < j_2 < \cdots < j_t\}$, $(1 \leqq t \leqq N, 1 \leqq j \leqq \alpha_t)$, we set $A^{(t)}_{j,k} = j_k$, and $A^{(t)}_{Min} = \{A^{(t)}_{1,1}, A^{(t)}_{2,1}, \cdots, A^{(t)}_{t,1}\}$.*

**Example**

$$\begin{aligned}
\mathcal{O}_1(z) &= T_1(z), && (4.5)\\
\mathcal{O}_2(z_1,z_2) &= g_{1,1}(z_2/z_1)T_1(z_1)T_1(z_2) - c\delta(x^2z_2/z_1)T_2(x^{-1}z_1), && (4.6)\\
\mathcal{O}_3(z_1,z_2,z_3) &= g_{11}(z_2/z_1)g_{1,1}(z_3/z_1)g_{1,1}(z_3/z_2)T_1(z_1)T_1(z_2)T_1(z_3)\\
&- cg_{1,2}(x^{-1}z_2/z_1)T_1(z_1)\delta(x^2z_3/z_2)T_2(x^{-1}z_2)\\
&- cg_{1,2}(x^{-1}z_1/z_2)T_1(z_2)\delta(x^2z_3/z_1)T_2(x^{-1}z_1)\\
&- cg_{1,2}(x^{-1}z_1/z_3)T_1(z_3)\delta(x^2z_2/z_1)T_2(x^{-1}z_1)\\
&+ c^2\Delta(x^3)(\delta(x^2z_2/z_1)\delta(x^2z_1/z_3) + \delta(x^2z_1/z_2)\delta(x^2z_3/z_1))T_3(z_1). && (4.7)
\end{aligned}$$

Let us set the auxiliary function $s(z) = s(1/z)$ by

$$s(z) = \frac{(z;x^{2s})_\infty (x^{2s-2r}z;x^{2s})_\infty}{(x^{2s-2}z;x^{2s})_\infty (x^{-2r+2}z;x^{2s})_\infty} \times \frac{(1/z;x^{2s})_\infty (x^{2s-2r}/z;x^{2s})_\infty}{(x^{2s-2}/z;x^{2s})_\infty (x^{-2r+2}/z;x^{2s})_\infty}. \tag{4.8}$$

**Definition 4.2** *For $\mathrm{Re}(s) > 0$ and $\mathrm{Re}(r) < 0$, we define a family of the operators $\mathcal{I}_n$, $(n = 1, 2, \cdots)$ by*

$$\mathcal{I}_n = \left[ \prod_{1\leqq j<k\leqq n} s(z_k/z_j)\mathcal{O}_n(z_1,\cdots,z_n) \right]_{1,z_1,\cdots,z_n}. \tag{4.9}$$

*For generic* $\mathrm{Re}(s) > 0$ *and* $r \in \mathbb{C}$*, the definition of* $\mathcal{I}_n$ *should be understood as analytic continuation. We call the operator* $\mathcal{I}_n$ *the Local Integral Motion for the deformed* $W$*-algebra* $W_{q,t}(\widehat{sl_N})$*.*

**Proposition 4.1** *The operator* $\mathcal{O}_n(z_1, z_2, \cdots, z_n)$ *is* $S_n$*-invariant in "weakly sense".*

$$\prod_{1 \leqq j<k \leqq n} s(z_k/z_j)\mathcal{O}_n(z_1, z_2, \cdots, z_n) = \prod_{1 \leqq j<k \leqq n} s(z_{\sigma(k)}/z_{\sigma(j)})\mathcal{O}_n(z_{\sigma(1)}, z_{\sigma(2)}, \cdots, z_{\sigma(n)}), \ (\sigma \in S_n). \tag{4.10}$$

The following is one of **Main Results**.

**Theorem 4.2** *The Local Integrals of Motion* $\mathcal{I}_n$ $(n = 1, 2, \cdots)$ *commute with each other.*

$$[\mathcal{I}_m, \mathcal{I}_n] = 0, \quad (m, n = 1, 2, \cdots). \tag{4.11}$$

By using $S_n$-invariance of $\mathcal{O}_n(z_1, z_2, \cdots, z_n)$ in "weakly sense", the above theorem is reduced to the following theta function identity, which is shown by induction [6].

$$\begin{aligned} &\sum_{\substack{J \subset \{1,2,\cdots,n+m\} \\ |J|=n}} \prod_{j \in J} \prod_{k \notin J} \frac{[u_k - u_j + 1]_s [u_k - u_j + r - 1]_s}{[u_k - u_j]_s [u_k - u_j + r]_s}, \\ = &\sum_{\substack{J^c \subset \{1,2,\cdots,n+m\} \\ |J^c|=m}} \prod_{j \in J^c} \prod_{k \notin J^c} \frac{[u_k - u_j + 1]_s [u_k - u_j + r - 1]_s}{[u_k - u_j]_s [u_k - u_j + r]_s}. \end{aligned} \tag{4.12}$$

**Conjecture 4.3** *The Local Integrals of Motion are* $\eta$*-invariant.*

$$\eta(\mathcal{I}_n) = \mathcal{I}_n, \quad (n = 1, 2, \cdots). \tag{4.13}$$

We have checked for small $n$. We have already shown $\eta$-invariance conjecture, $\eta(\mathcal{I}_n) = \mathcal{I}_n$, for the deformed Virasoro algebra case, $Vir_{q,t} = W_{q,t}(\widehat{sl_2})$ [8].

# 5 Nonlocal Integrals of Motion

In this section we give explicit formulae of the Nonlocal Integrals of Motion. Let us set the theta function $\vartheta(u^{(1)}|u^{(2)}|\cdots|u^{(N)})$ by following conditions.

$$\vartheta(u^{(1)}|\cdots|u^{(t)} + r|\cdots|u^{(N)}) = \vartheta(u^{(1)}|\cdots|u^{(t)}|\cdots|u^{(N)}), \ (1 \leqq t \leqq N) \tag{5.1}$$

$$\begin{aligned}
&\vartheta(u^{(1)}|\cdots|u^{(t)}+r\tau|\cdots|u^{(N)}) \\
= \ & e^{-2\pi i\tau-\frac{2\pi i}{r}(u_{t-1}-2u_t+u_{t+1}+\sqrt{r(r-1)}P_{\alpha_t})}\vartheta(u^{(1)}|\cdots|u^{(t)}|\cdots|u^{(N)}), \quad (1 \leqq t \leqq N), \qquad (5.2) \\
&\vartheta(u^{(1)}+k|\cdots|u^{(N)}+k) = \vartheta(u^{(1)}|\cdots|u^{(N)}), \quad (k \in \mathbb{C}), \qquad (5.3) \\
&\eta(\vartheta(u^{(1)}|\cdots|u^{(N)})) = \vartheta(u^{(N)}|u^{(1)}|\cdots|u^{(N-1)}). \qquad (5.4)
\end{aligned}$$

**Example** For $N=2$ case, we have

$$\begin{aligned}
\vartheta(u_1|u_2) &= [u_1-u_2-\sqrt{r(r-1)}P_{\alpha_1}+\alpha]_r[u_1-u_2-\alpha]_r \\
&+ [u_1-u_2-\sqrt{r(r-1)}P_{\alpha_1}-\alpha]_r[u_1-u_2+\alpha]_r, \quad (\alpha \in \mathbb{C}). \qquad (5.5)
\end{aligned}$$

**Definition 5.1** *For* $\mathrm{Re}(r) \neq 0$ *and* $0 < \mathrm{Re}(s) < 2$*, we define a family of operators* $\mathcal{G}_m, \ (m=1,2,\cdots)$ *by*

$$\begin{aligned}
\mathcal{G}_m &= \prod_{t=1}^{N}\prod_{j=1}^{m}\oint_C \frac{dz_j^{(t)}}{2\pi\sqrt{-1}z_j^{(t)}} F_1(z_1^{(1)})\cdots F_1(z_m^{(1)})F_2(z_1^{(2)})\cdots F_2(z_m^{(2)})\cdots F_N(z_1^{(N)})\cdots F_N(z_m^{(N)}) \\
&\times \frac{\displaystyle\prod_{t=1}^{N}\prod_{1\leqq i<j\leqq m}\left[u_i^{(t)}-u_j^{(t)}\right]_r\left[u_j^{(t)}-u_i^{(t)}-1\right]_r}{\displaystyle\prod_{t=1}^{N-1}\prod_{i,j=1}^{m}\left[u_i^{(t)}-u_j^{(t+1)}+1-\frac{s}{N}\right]_r\prod_{i,j=1}^{m}\left[u_i^{(1)}-u_j^{(N)}+\frac{s}{N}\right]_r} \\
&\times \vartheta\left(\sum_{j=1}^{m}u_j^{(1)}\middle|\sum_{j=1}^{m}u_j^{(2)}\middle|\cdots\middle|\sum_{j=1}^{m}u_j^{(N)}\right). \qquad (5.6)
\end{aligned}$$

*Here the integral contour* $C$ *is given by*

$$\begin{aligned}
&|x^{\frac{2s}{N}}z_j^{(t+1)}| < |z_i^{(t)}| < |x^{-2+\frac{2s}{N}}z_j^{(t+1)}|, \quad (1 \leqq t \leqq N-1, 1 \leqq i,j \leqq m), \qquad (5.7) \\
&|x^{2-\frac{2s}{N}}z_j^{(1)}| < |z_i^{(N)}| < |x^{-\frac{2s}{N}}z_j^{(1)}|, \quad (1 \leqq i,j \leqq m). \qquad (5.8)
\end{aligned}$$

*For generic* $s \in \mathbb{C}$*, the definition of* $\mathcal{G}_n$ *should be understood as analytic continuation. We call the operator* $\mathcal{G}_n$ *the Nonlocal Integrals of Motion for the deformed* $W$*-algebra* $W_{q,t}(\widehat{sl_N})$*.*

**Example** For $N=2$ and $m=1$ case, we have

$$\mathcal{G}_1 = \int\int_C \frac{dz_1}{2\pi\sqrt{-1}z_1}\frac{dz_2}{2\pi\sqrt{-1}z_2}F_1(z_1)F_0(z_2)\frac{\vartheta(u_1|u_2)}{[u_1-u_2+\frac{s}{2}]_r[u_1-u_2-\frac{s}{2}+1]_r}. \qquad (5.9)$$

Here $C$ is given by $|x^s z_2| < |z_1| < |x^{-2+s} z_2|$.

The following is one of **Main Results**.

**Theorem 5.1** *The Nonlocal Integrals of Motion* $\mathcal{G}_n$, $(n = 1, 2, \cdots)$ *commute with each other.*

$$[\mathcal{G}_m, \mathcal{G}_n] = 0, \quad (m, n = 1, 2, \cdots). \tag{5.10}$$

By using commutation relations of the screening currents $F_j(z)$, the above theorem is reduced to the following theta function identity, which is shown by induction [6].

$$\begin{aligned}
& \sum_{\sigma_1 \in S_{m+n}} \sum_{\sigma_2 \in S_{m+n}} \cdots \sum_{\sigma_N \in S_{m+n}} \vartheta_\alpha \left( \sum_{j=1}^{m} u^{(1)}_{\sigma_1(j)} \middle| \sum_{j=1}^{m} u^{(2)}_{\sigma_2(j)} \middle| \cdots \middle| \sum_{j=1}^{m} u^{(N)}_{\sigma_N(j)} \right) \\
\times\ & \vartheta_\beta \left( \sum_{j=m+1}^{m+n} u^{(1)}_{\sigma_1(j)} \middle| \sum_{j=m+1}^{m+n} u^{(2)}_{\sigma_2(j)} \middle| \cdots \middle| \sum_{j=m+1}^{m+n} u^{(N)}_{\sigma_N(j)} \right) \\
\times\ & \prod_{t=1}^{N} \prod_{i=1}^{m} \prod_{j=m+1}^{m+n} \frac{\left[ u^{(t)}_{\sigma_t(i)} - u^{(t+1)}_{\sigma_{t+1}(j)} - \frac{s}{N} \right]_r \left[ u^{(t)}_{\sigma_t(j)} - u^{(t+1)}_{\sigma_{t+1}(i)} + 1 - \frac{s}{N} \right]_r}{\left[ u^{(t)}_{\sigma_t(i)} - u^{(t)}_{\sigma_t(j)} \right]_r \left[ u^{(t)}_{\sigma_t(j)} - u^{(t)}_{\sigma_t(i)} - 1 \right]_r} \\
=\ & \sum_{\sigma_1 \in S_{m+n}} \sum_{\sigma_2 \in S_{m+n}} \cdots \sum_{\sigma_N \in S_{m+n}} \vartheta_\beta \left( \sum_{j=1}^{n} u^{(1)}_{\sigma_1(j)} \middle| \sum_{j=1}^{n} u^{(2)}_{\sigma_2(j)} \middle| \cdots \middle| \sum_{j=1}^{n} u^{(N)}_{\sigma_N(j)} \right) \\
\times\ & \vartheta_\alpha \left( \sum_{j=n+1}^{m+n} u^{(1)}_{\sigma_1(j)} \middle| \sum_{j=n+1}^{m+n} u^{(2)}_{\sigma_2(j)} \middle| \cdots \middle| \sum_{j=n+1}^{m+n} u^{(N)}_{\sigma_N(j)} \right) \\
\times\ & \prod_{t=1}^{N} \prod_{i=1}^{n} \prod_{j=n+1}^{m+n} \frac{\left[ u^{(t)}_{\sigma_t(i)} - u^{(t+1)}_{\sigma_{t+1}(j)} - \frac{s}{N} \right]_r \left[ u^{(t)}_{\sigma_t(j)} - u^{(t+1)}_{\sigma_{t+1}(i)} + 1 - \frac{s}{N} \right]_r}{\left[ u^{(t)}_{\sigma_t(i)} - u^{(t)}_{\sigma_t(j)} \right]_r \left[ u^{(t)}_{\sigma_t(j)} - u^{(t)}_{\sigma_t(i)} - 1 \right]_r},
\end{aligned} \tag{5.11}$$

where $\vartheta_\alpha(u^{(1)}|u^{(2)}|\cdots|u^{(N)})$ and $\vartheta_\beta(u^{(1)}|u^{(2)}|\cdots|u^{(N)})$ are not necesarry the same theta functions.

**Theorem 5.2** *The Nonlocal Integrals of Motion are* $\eta$*-invariant.*

$$\eta(\mathcal{G}_m) = \mathcal{G}_m, \quad (m = 1, 2, \cdots). \tag{5.12}$$

**Conjecture 5.3** *The Local Integrals of Motion* $\mathcal{I}_n$, $(n = 1, 2, \cdots)$ *and Nonlocal Integrals of Motion* $\mathcal{G}_m$, $(m = 1, 2, \cdots)$ *commute with each other.*

$$[\mathcal{I}_n, \mathcal{G}_m] = 0, \quad (m, n = 1, 2, \cdots). \tag{5.13}$$

We have already shown the commutation relations $[\mathcal{I}_n, \mathcal{G}_m] = 0, (m, n = 1, 2, \cdots)$ for the deformed Virasoro algebra case, $Vir_{q,t} = W_{q,t}(\widehat{sl_2})$ [8]. We have already shown the commutation relations $[\mathcal{I}_1, \mathcal{G}_m] = 0$, $(m = 1, 2, \cdots)$ for general $W_{q,t}(\widehat{sl_N})$ case. If we assume $\eta$-invariance of the Local Integrals of Motion $\eta(\mathcal{I}_n) = \mathcal{I}_n$, the commutation relation $[\mathcal{I}_n, \mathcal{G}_m] = 0$, $(m, n = 1, 2, \cdots)$ for general $W_{q,t}(\widehat{sl_N})$, can be shown by simple argument.

**Acknowledgements.** We would like to thank Professor M.Jimbo for his interest to this work. B.F.is partly supported by Grant RFBR (05-01-01007), SS (2044.2003.2), and RFBR-JSPS (05-01-02934). T.K. is partly supported by Grant-in Aid for Young Scientist **B** (18740092) from JSPS. J.S. is partly supported by Grant-in Aid for Scientific Research **C** (16540183) from JSPS.

# References

[1] V.Bazhanov, S.Lukyanov, Al.Zamolodchikov : Integral Structure of Conformal Field Theory, Quantum KdV Theory and Thermodynamic Bethe Ansatz *Commun.Math.Phys.***177**, no.2, 381-398, (1996).

[2] V.Bazhanov, A.Hibberd, S.Khoroshkin : Integrable Structure of $W_3$ Conformal Field Theory, *Nucl.Phys.***B622** 475-547,(2002).

[3] J.Shiraishi, H.Kubo, H.Awata, S.Odake: A Quantum Deformation of the Virasoro Algebra and the Macdonald Symmetric Functions, *Lett.Math.Phys.***38**, 647-666, (1996).

[4] H.Awata, H.Kubo, S.Odake, J.Shiraishi : Quantum $W_N$ Algebras and Macdonald Polynomials, *Commun.Math.Phys.***179**, 401-416, (1996).

[5] S.Odake : Comments on the Deformed $W_N$ Algebra, APCTP-Nankai Joint Symposium on "Lattice Statistics and Mathematical Physics 2001", Tianjin China.

[6] B.Feigin and A.Odesskii : A Family of Elliptic Algebras, *Internat.Math.Res.Notices* no.11, 531-539, (1997).

[7] B.Feigin, M.Jimbo, T.Miwa, A.Odesskii, Ya.Pugai : Algebra of Screening Operators for the Deformed $W_n$ Algebra, *Commun. Math. Phys.* **191**, 501-541, (1998).

[8] B.Feigin, T.Kojima, J.Shiraishi, H.Watanabe : The Integrals of Motion for the Deformed Virasoro Algebra, Preprint, (2006),[arXiv:0705.0427].

*Proceedings for Representation Theory* 2006, Atami, Japan, p.102-114 (2006).

[ISBN4-9902328-2-8]